arXiv:1302.6771v1 [q-bio.PE] 27 Feb 2013

A two-fold advantage of sex

Su-Chan Park[*] and Joachim Krug[†]

February 28, 2013

[*] Department of Physics, The Catholic University of Korea, Bucheon 420-743, Republic of Korea

[†] Institut für Theoretische Physik, Universität zu Köln, 50937 Köln, Germany

Running Head: A two-fold advantage of sex

Key Words: advantage of recombination, clonal interference, speed of adaptation

Corresponding Author:

Su-Chan Park

Department of Physics
The Catholic University of Korea
43 Jibong-ro, Wonmi-gu,
Bucheon 420-743, Republic of Korea
Tel : +82-2-2164-4524
e-mail : spark0@catholic.ac.kr

Abstract

The adaptation of large asexual populations is hampered by the competition between independently arising beneficial mutations in different individuals, which is known as clonal interference. In classic work, Fisher and Muller proposed that recombination provides an evolutionary advantage in large populations by alleviating this competition. Based on recent progress in quantifying the speed of adaptation in asexual populations undergoing clonal interference, we present a detailed analysis of the Fisher-Muller mechanism for a model genome consisting of two loci with an infinite number of beneficial alleles each and multiplicative (non-epistatic) fitness effects. We solve the deterministic, infinite population dynamics exactly and show that, for a particular, natural mutation scheme, the speed of adaptation in sexuals is twice as large as in asexuals. This result is argued to hold for any nonzero value of the rate of recombination. Guided by the infinite population result and by previous work on asexual adaptation, we postulate an expression for the speed of adaptation in finite sexual populations that agrees with numerical simulations over a wide range of population sizes and recombination rates. The ratio of the sexual to asexual adaptation speed is a function of population size that increases in the clonal interference regime and approaches 2 for extremely large populations. The simulations also show that recombination leads to a strong equalization of the number of fixed mutations in the two loci. The generalization of the model to an arbitrary number L of loci is briefly discussed. For a particular communal recombination scheme, the ratio of the sexual to asexual adaptation speed is approximately equal to L in large populations.

The evolutionary advantage of sex remains one of the most intriguing puzzles in evolutionary biology (KONDRASHOV 1993; DE VISSER and ELENA 2007; OTTO 2009). Many hypotheses have been suggested explaining why sexual reproduction is widespread in nature despite apparent disadvantages such as the two-fold cost of sex (MAYNARD SMITH 1978). Well-known examples are the deterministic mutation hypothesis (KONDRASHOV 1988), the Fisher-Muller mechanism (FISHER 1930; MULLER 1932; CROW and KIMURA 1965) and Muller's ratchet (MULLER 1964; FELSENSTEIN 1974), to name only a few. These three hypotheses are applicable when the fitness landscape in question has certain specific features. Specifically, the deterministic mutation hypothesis requires deleterious mutations to be synergistically epistatic, while the Fisher-Muller (FM) mechanism as well as Muller's ratchet can explain the advantage of sex if epistasis is negligible.

Theoretical analyses of the effect of epistasis on the speed of Muller's ratchet have concluded that it practically stops operating when epistasis is synergistic (CHARLESWORTH *et al.* 1993; KONDRASHOV 1994; JAIN 2008). Furthermore, recent experimental analyses of empirical fitness landscapes seem to indicate that a particularly strong form of epistasis termed sign epistasis (WEINREICH *et al.* 2005) is the rule rather than an exception (WEINREICH *et al.* 2006; DE VISSER *et al.* 2009; FRANKE *et al.* 2011; SZENDRO *et al.* 2013). Sign epistasis generally implies that the fitness landscape is rugged. On a rugged fitness landscape sex can be detrimental, even without taking into account the two-fold cost of sex, in that sexual populations, unlike the corresponding asexual populations, cannot escape from local fitness peaks (CROW and KIMURA 1965; ESHEL and FELDMAN 1970; DE VISSER *et al.* 2009; PARK and KRUG 2011).

Although research on empirical fitness landscapes has been growing substantially in recent years, it is still practically infeasible to reliably determine genotypic fitness on a genome-wide scale [but see KOUYOS *et al.* (2012)]. Because of the small sizes of most empirical fitness landscapes that have so far been constructed experimentally, the implications of the ubiquity of sign epistasis observed in these studies for long term evolution remain unclear. At the same time experimental evidence in favor of the FM mechanism has accumulated (Colegrave 2002; Cooper 2007). For these reasons further quantitative analysis of the advantage of sex in the *absence* of epistasis remains a worthwhile endeavor, and we will pursue this approach in the present contribution.

The essence of the FM mechanism is the competition between independently arising beneficial mutations, termed clonal interference, which slows down the adaptation of large asexual populations (Gerrish and Lenski 1998; Miralles *et al.* 1999; Wilke 2004; Kim and Orr 2005; Park and Krug 2007; Fogle *et al.* 2008; Sniegowski and Gerrish 2010; Schiffels *et al.* 2011). The concept of clonal interference has played an important role in intepreting the behavior observed in laboratory selection experiments (Lenski *et al.* 1991; Lenski and Travisano 1994; Barrick *et al.* 2009), and has also been invoked in explaining the population-size dependence of evolutionary predictability in rugged fitness landscapes (Jain *et al.* 2011; Szendro *et al.* 2013). Although in its original formulation clonal interference theory neglects the occurrence of secondary beneficial mutations within a growing clone (Gerrish and Lenski 1998; Gerrish 2001), in general the coexistence of multiple beneficial mutations cannot be neglected in large populations (Park and Krug 2007). In the following we will therefore use the term clonal interference in a wider sense than originally conceived, in that two clones with different numbers of beneficial mutations can compete with each other for fixation.

Much recent theoretical work has focused on obtaining accurate quantitative estimates of the speed of adaptation in the presence of clonal interference for the simple situation of an unlimited supply of beneficial mutations that act independently on fitness, without epistatic interactions [see Park *et al.* (2010) for review]. It turns out that the population dynamics in this regime is well described by a traveling wave moving at constant speed along a one-dimensional fitness space. The traveling wave picture was first established for the case when the beneficial selection coefficient is the same for all mutations (Tsimring *et al.* 1996; Rouzine *et al.* 2003; Desai and Fisher 2007; Brunet *et al.* 2008; Rouzine *et al.* 2008) and recently extended to the more realistic case of selection coefficients drawn from a continuous effect size distribution (GOOD *et al.* 2012; FISHER 2013).

In natural populations it is unlikely that the traveling wave picture persists forever. Apart from the assumed absence of epistatic interactions, there are two main features that lead to a breakdown of this picture in long-term evolution. First, a fluctuating environment generally makes the fitness landscape change with time. In a time-dependent situation it is problematic to compare fitnesses of two individuals living on different landscapes and the notion of absolute fitness has to be replaced by fitness flux (MUSTONEN and LÄSSIG 2010). Second, even if the fitness landscape remains constant for a very long time, the indefinite supply of beneficial mutations appearing at constant rate cannot be a good approximation in the real world. For example, in long-term evolution experiments the speed of adaptation usually slows down (LENSKI and TRAVISANO 1994; BARRICK *et al.* 2009), which is attributed to the decreasing supply of beneficial mutations. In this context, the house-of-cards model, in which fitness values are assigned randomly to genotypes, could provide a more realistic description (KINGMAN 1978; PARK and KRUG 2008). In the framework of this model one cannot however explain the advantage of sex, because recombination is no different from mutation.

Although the non-epistatic model with an infinite supply of beneficial mutations is of limited validity, it can provide a reasonable approximation when a population undergoes a severe environmental change, as is often the case at the beginning of an evolution experiment. At the same time this setting is conceptually simple and allows for detailed (if approximate) mathematical analysis. In the present paper, we therefore build upon the recent line of work on asexual populations undergoing clonal interference and add to it a minimal yet realistic recombination scheme.

Specifically, we consider a facultatively sexual population model with two genetic loci, each of which can acquire infinitely many beneficial mutations. We assume that epistasis is absent both between and within loci. We first analyze the infinite population dynamics, obtaining exact expressions for the speed of adaptation in the limiting cases of zero and maximal recombination rates (asexuals vs. obligate sexuals). When the selection coefficient of beneficial mutations is the same at both loci and at most one mutation may occur per generation and individual, the speed of adaptation for obligate sexuals is twice that of asexuals, a result that we argue holds for any positive recombination rate. Based on this observation and on previous results for asexual populations, we propose a simple expression for the speed of adaptation in finite sexual populations that is in good agreement with simulations and predicts a roughly 'two-fold advantage of sex' for large populations and moderate recombination rates. In DISCUSSION we describe a possible extension of the model to more than two loci, and place our work into the context of related studies.

MODELS

We consider a (sexual or asexual) population of haploid individuals in discrete generations. The population size is denoted by N and assumed to be constant. As a reproduction scheme we employ the Wright-Fisher model (Fisher 1930; Wright 1931), the prototypical model of discrete, non-overlapping generations. Since our main concern is how recombination affects the speed of adaptation, we assume that all mutations are beneficial. This naturally leads us to study evolution in the framework of the infinite-sites model (Kimura 1969); otherwise back mutations of beneficial mutations, which are deleterious by definition, should appear with nonzero probability. Furthermore, we assume no epistasis among mutations, which will be reflected by the multiplicative fitness assignement (see below). As a minimal model with the above properties, we study an evolving population with only two loci under selection. Each locus is assumed to have infinitely many sites. We assume an initially homogenous population and the fitness of the initial genome, or wild-type, is set to unity.

In line with the assumption of multiplicative fitness effects, the fitness of an individual that has n_i mutations at locus i compared to the wild-type is $\exp(n_1 s_1 + n_2 s_2)$. Without loss of generality we take $s_2 \geq s_1$. Note that two genotypes with the same number of mutations at each locus are not necessarily the same though both have the same fitness $\exp(s_1 n_1 + s_2 n_2)$.

Since we are only interested in how fast mean fitness increases and not in the genealogy, all genotypes with the same number of mutations at each locus will be treated as if they were the same.

The population evolves in the following way. Let $f_t(n_1, n_2)$ denote the frequency of all genotypes with n_1 (n_2) mutations at the first (second) locus in generation t ($t = 0, 1, 2, \ldots$). Initially, the population is homogeneous with $f_t(n_1, n_2) = \delta_{n_1,0}\delta_{n_2,0}$, where $\delta_{x,y}$ is the Kronecker delta symbol which is 1 (0) if $x = y$ ($x \neq y$). By selection, the frequency at generation $t + 1$ on average will change to be

$$f_t^s(n_1, n_2) = \frac{e^{s_1 n_1 + s_2 n_2}}{\bar{w}_t} f_t(n_1, n_2), \tag{1}$$

where

$$\bar{w}_t \equiv \sum_{n_1, n_2} e^{s_1 n_1 + s_2 n_2} f_t(n_1, n_2) \tag{2}$$

is the average fitness of the population at generation t.

Mutation can also change the frequency of genotypes. Let $g_0(m_1, m_2)$ denote the probability that mutation changes m_i sites of the i-th locus ($i = 1, 2$) of an offspring ($m_1, m_2 \geq 0$). Here we implicitly assume that the mutation probability is not affected by the genetic background. Note that the probability of having no mutation is included in g_0 for convenience. The frequency change due to both selection and mutation is

$$\begin{aligned} f_t^\mu(n_1, n_2) &= \sum_{m_1, m_2 \geq 0} g_0(n_1 - m_1, n_2 - m_2) f_t^s(m_1, m_2) \\ &= \sum_{m_1, m_2 \geq 0} g_0(n_1 - m_1, n_2 - m_2) \frac{e^{s_1 m_1 + s_2 m_2}}{\bar{w}_t} f_t(m_1, m_2), \end{aligned} \tag{3}$$

where $g_0(x, y)$ with at least one negative argument should be understood to be 0. We further assume that mutation does not have any preference for a certain locus, that is, $g_0(m_1, m_2) = g_0(m_2, m_1)$ for any pair of m_1 and m_2.

After selection and mutation, two randomly chosen parents mate and beget an offspring. Let $R(n_1, n_2 | k_1, k_2; l_1, l_2)$ denote the probability that the resulting progeny of two individuals with respective genotypes (k_1, k_2) and (l_1, l_2) has the genotype (n_1, n_2). To be specific, we set $(0 \leq r \leq 1)$

$$R(n_1, n_2|k_1, k_2; l_1, l_2) = \frac{1-r}{2}\left(\delta_{n_1k_1}\delta_{n_2k_2} + \delta_{n_1l_1}\delta_{n_2l_2}\right) + \frac{r}{2}\left(\delta_{n_1k_1}\delta_{n_2l_2} + \delta_{n_1l_1}\delta_{n_2k_2}\right), \tag{4}$$

which means that with probability $1-r$ the two loci of the offspring in question are inherited solely from a single parent and with probability r the offspring inherits one locus from one parent and the other from the other parent. Note that the case with $r = \frac{1}{2}$ is generally referred to as the free recombination, while $r = 1$ corresponds to obligate sexuals.

Since the probability that the randomly chosen parents have genotypes (k_1, k_2) and (l_1, l_2) is $f_t^\mu(k_1, k_2) f_t^\mu(l_1, l_2)$, the mean frequency after selection, mutation, and recombination is

$$\begin{aligned} f_t^r(n_1, n_2) &= \sum_{k_1,k_2,l_1,l_2} R(n_1, n_2|k_1, k_2; l_1, l_2) f_t^\mu(k_1, k_2) f_t^\mu(l_1, l_2) \\ &= (1-r) f_t^\mu(n_1, n_2) + r f_t^{(1)}(n_1) f_t^{(2)}(n_2) \end{aligned} \tag{5}$$

where

$$f_t^{(1)}(n_1) = \sum_{n_2} f_t^\mu(n_1, n_2), \quad f_t^{(2)}(n_2) = \sum_{n_1} f_t^\mu(n_1, n_2), \tag{6}$$

are marginal frequency distributions of genotypes after the selection and mutation steps with n_1 mutations at locus 1 and n_2 mutations at locus 2, respectively.

Finally, the actual population distribution at generation $t+1$ is determined by multinomial sampling using $f_t^r(n_1, n_2)$ in Equation 5 with the restriction that the population size is N. For simulations, we employ the algorithm explained by PARK and KRUG (2007) [see also PARK *et al.* (2010) for simulations of extremely large populations].

The speed of adaptation, or shortly speed, is defined as the rate of increase of the log mean fitness,

$$v \equiv \lim_{t\to\infty} \frac{\langle \ln \bar{w}_t \rangle}{t}, \tag{7}$$

where $\langle \ldots \rangle$ denotes an average over independent realizations of evolution with the same parameters. In the following, we mainly focus on the dependence of speed on parameters such as the mutation probability per generation, the selection coefficient of a single mutation, and the recombination probability r.

RESULTS

Infinite populations: Although the infinite population limit cannot be reached in real biological populations for the model we are studying (Park *et al.* 2010), it does provide some insight into the adaptation dynamics of finite populations. Furthermore, the deterministic nature of the infinite population dynamics renders an analytic approach feasible.

When the population size is infinite, $f_{t+1}(n_1, n_2)$ is equal to $f_t^r(n_1, n_2)$ by the law of large numbers. In APPENDIX A, we derive the mean fitness from the recursion relation for the frequencies in two special cases, $r = 0$ and $r = 1$. From Equations A7 and A11, we deduce the speed of adaptation as

$$v_a \equiv v(r = 0, N = \infty) = \lim_{t\to\infty} \frac{\ln G_0(e^{s_1 t}, e^{s_2 t})}{t}, \tag{8}$$

$$v_s \equiv v(r = 1, N = \infty) = \lim_{t\to\infty} \frac{\ln \widetilde{G}_0(e^{s_1 t}) + \ln \widetilde{G}_0(e^{s_2 t})}{t}, \tag{9}$$

where subscripts a and s stand for asexuals and obligate sexuals, respectively, G_0 is the generating function for g_0, defined as

$$G_0(z_1, z_2) \equiv \sum_{k_1,k_2} z_1^{k_1} z_2^{k_2} g_0(k_1, k_2), \tag{10}$$

and $\widetilde{G}_0(z) = G_0(z, 1) = G_0(1, z)$ is the generating function for the marginal distribution $\tilde{g}_0(k) = \sum_m g_0(k, m) = \sum_m g_0(m, k)$ [see also Equation A5]. Note that we are using the symmetry $g_0(k_1, k_2) = g_0(k_2, k_1)$ introduced earlier, but the generalization to asymmetric g_0 is straightforward.

Since the arguments of G_0 in Equation 8 and of $\widetilde{G}_0$ in Equation 9 increase exponentially, the speed is fully determined by the largest possible fitness effect due to a single mutation event. Thus,

$$\begin{aligned} v_a &= \mathrm{Max}\{n_1 s_1 + n_2 s_2 | g_0(n_1, n_2) \neq 0\}, && (11) \\ v_s &= M(s_1 + s_2), && (12) \end{aligned}$$

where M is the largest possible number of mutated sites at *one locus* by a single mutation event,

$$M = \mathrm{Max}\{n | \exists n' \text{ such that } g_0(n, n') = g_0(n', n) \neq 0\}. \qquad (13)$$

Since, by definition, M is the maximum of all possible n_1 and n_2 with $g(n_1, n_2) \neq 0$, v_s cannot be smaller than v_a. Thus, sex is at least not detrimental, though it may have no effect depending on the form of g_0. For example, if single mutations occur with probability U and double mutations involving both loci with probability U^2, corresponding to

$$g_0(k, m) = (1 - U - U^2)\delta_{k+m,0} + \frac{U}{2}\delta_{k+m,1} + U^2 \delta_{k,1}\delta_{m,1}, \qquad (14)$$

then $v_s = v_a = s_1 + s_2$. On the other hand, if double mutations are forbidden and

$$g_0(k, m) = (1 - U)\delta_{k+m,0} + \frac{U}{2}\delta_{k+m,1}, \qquad (15)$$

we have $v_s = s_1 + s_2 > v_a = s_2$ (recall that we assume $s_2 \geq s_1$). Hence the effect of sex significantly depends on the form of g_0 in the infinite population limit. If $s_2 > s_1$ (strict inequality) and if g_0 is as in Equation 15, beneficial mutations occurring on locus 1 do not contribute to the speed of an infinite asexual population. This can be understood in the framework of clonal interference as the 'wasting' of weaker beneficial mutations by the competition with stronger mutations. On this account, we will later assume $s_1 = s_2$ when we study the adaptation dynamics of finite populations in order to maximize the effect of both loci in the asexual case. Note that for $s_1 = s_2$ the mutation scheme in Equation 15 implies $v_s = 2v_a$, a two-fold advantage of sex.

Although we only calculated the speed exactly for the cases $r = 0$ and $r = 1$, we now argue that the asymptotic speed does not depend on r provided $r > 0$. Let $\ell_1(t)$ ($\ell_2(t)$) denote the maximum number of mutations at locus 1 (locus 2) accumulated up to generation t:

$$\ell_1(t) \equiv \max\left\{ n \left| \sum_m f_t(n,m) \neq 0 \right. \right\}, \ \ell_2(t) \equiv \max\left\{ m \left| \sum_n f_t(n,m) \neq 0 \right. \right\}. \tag{16}$$

This definition can be used for finite populations as well, and is closely related to the *lead* of the fitness distribution considered in the traveling wave approach to asexual adaptation (DESAI and FISHER 2007; PARK *et al.* 2010; FISHER 2013). Since the population size is infinite, the (marginal) frequency of genomes with $\ell_i(t) + M$ mutations at the ith locus at generation $t+1$, where M was defined in Equation 13, is nonzero, though it can be extremely small. Thus, by recombination, a nonzero fraction of individuals will have $2M + \ell_1(t) + \ell_2(t)$ mutations in total. Combined with the initial condition $\ell_1(0) = \ell_2(0) = 0$, the detectable largest fitness at generation t is then $M(s_1 + s_2)t$ for all $t \geq 0$. Now assume that the speed $v(r, N = \infty)$ for $0 < r < 1$ is strictly smaller than $M(s_1 + s_2)$. Then, with time t, the ratio of the detectable largest fitness to the mean fitness increases as $\exp((M(s_1 + s_2) - v)t)$. Thus, at some point, $f_t(Mt, Mt)$ jumps abruptly up to a significant number in a single generation, which eventually leads to an abrupt increase of $\bar{w}_t$ up to the order of $\exp(M(s_1 + s_2)t)$. Hence, in the long run, the speed becomes $M(s_1 + s_2)$ for any $r > 0$.

In the above discussion, we argued that the speed does not depend on r once r is nonzero. On the other hand, if r is very small, the whole population behaves almost like an asexual population for quite some time. Hence, the abrupt jump of fitness mentioned above should be observable. To see this phenomenon, we studied the deterministic evolution numerically, using the mutation scheme of Equation 15 with $U = 0.1$ and $s_1 = s_2 = 0.02$. In Fig. 1, we show how the mean fitness behaves with time for $r = 0$, $r = 10^{-9}$, and $r = 1$. Even for the minute recombination rate of $r = 10^{-9}$, the mean fitness closely follows the $r = 1$ curve, however with some oscillations. To eludicate the origin of this behavior we need to consider how the frequency distribution changes with time.

[Figure 1 about here.]

In the asexual case ($r = 0$) the frequency distribution over the number of mutations is well described by a Gaussian (Park *et al.* 2010). Furthermore, the frequency distribution of the obligately sexual population with $r = 1$ should also be well described by a Gaussian, because the generating function is just the product of two generating functions of asexual evolution (see Equation A8). However, for $0 < r \ll 1$, the Gaussian may not be a good approximation. In Fig. 2 we depict the time evolution of the frequency distribution for $r = 10^{-9}$. Clearly the frequency distribution cannot be approximated by a Gaussian traveling wave. Moreover, the shape of the distribution changes with time, which implies that there is no time-independent steady state. Rather, the distribution behaves like a 'breathing traveling wave' in that the behavior seen in Fig. 2 repeats periodically. The time when two peaks become comparable in Fig. 2 corresponds to the abrupt jump of mean fitness alluded to above. Further mathematical analysis of this phenomenon seems interesting, but we will not pursue it here because it is hardly observable in real, finite populations.

[Figure 2 about here.]

Finite populations: We mentioned above that many analytic approaches have been developed to find an expression for the speed of adaptation in large asexual populations (Rouzine *et al.* 2003; Desai and Fisher 2007; Brunet *et al.* 2008; Rouzine *et al.* 2008). Park *et al.* (2010) summarized these developments and compared simulation results with the proposed analytic expressions. The approximation of Rouzine *et al.* (2008) turned out to be quite accurate in a wide range of parameters. The only disadvantage of this approach is that the speed is obtained as an implicit function of N (see below). In this section, we will find a mathematical formula for the speed of adaptation in sexual populations, using both the suggested formula for asexuals and the results for the infinite population dynamics in the previous section.

For an infinite population, as we have seen above, the detailed form of the mutational probability distribution $g_0(k_1, k_2)$ affects the speed. However, for plausible values of the mutation rate and the selection coefficient such infinite population effects become observable only for unrealistically large populations. To be concrete, it was shown by Park *et al.* (2010) that population sizes of order $N \sim 10^{1500}$ are required to reach the infinite population regime for $s = 0.02$ and $U = 10^{-5}$. Hence, for a finite population of realistic size, the exact form of the mutation probability is not important provided the probability of having n mutations in one generation decreases as U^n with $U \ll 1$. In the following we therefore use Equation 15 and set $s_1 = s_2 = s$ for the reasons mentioned previously.

We begin with a discussion of the speed for asexual populations. As was illustrated by Park *et al.* (2010), the speed for the asexual version of our model ($r = 0$) is well approximated by the implicit equation

$$\ln N \approx \frac{v_a^{\mathrm{RBW}}}{2s^2}\left(\ln^2 \frac{v_a^{\mathrm{RBW}}}{eUs} + 1\right) - \ln \sqrt{\frac{s^3 U}{v_a^{\mathrm{RBW}} \ln(v_a^{\mathrm{RBW}}/(Us))}}, \tag{17}$$

where the subscript a in v_a refers to the asexual population, the superscript RBW refers to the authors of Rouzine *et al.* (2008), and $e \approx 2.718182$ is the base of the natural logarithm.

Since the approximation of the fitness distribution by a continuous traveling wave was used to derive Equation 17, it should not be surprising that the discrepancy between theory and simulation becomes relatively large when the size of population is small enough to realize the strong-selection weak-mutation (SSWM) regime, where the population is mostly monomorphic. Based on this observation, there is room for improvement of the approximation in an *ad-hoc* way as follows: First we note that the first (second) term in Equation 17 is dominant when the speed is high (low). Thus, when the population size is small, we can neglect the first term. Recalling that the speed in the SSWM regime is given by $v_a = 2NUs^2$ (Park *et al.* 2010), we modify Equation 17 as

$$\ln N \approx \frac{v_a}{2s^2}\left(\ln^2 \frac{v_a}{eUs} + 1\right) + \ln \frac{v_a}{2s^2 U} \tag{18}$$

which keeps the large speed behavior unchanged and enforces the SSWM result for small speeds. In Fig. 3 we show that Equation 18 provides a more accurate approximation to the speed obtained from simulations with $U = 10^{-6}$ and $s = 0.01$ than Equation 17.

[Figure 3 about here.]

Now we move on to the speed of sexual populations. At first, let us start from the case of $r = 1$ whose infinite population limit allows for an exact solution. As we show in APPENDIX A, the evolutionary dynamics of an infinite population with $r = 1$ can be viewed as the independent evolution of each locus with the marginal mutation probability $\tilde{g}_0(k)$. That is, we can divide the evolutionary dynamics into two independent asexual populations with reduced mutation probability and the speed of the sexual population is obtained by simply adding the speeds of these two virtual asexual populations. Within the mutation scheme given by Equation 15 with selection coefficients $s_1 = s_2 = s$ this implies that

$$v_s\left(r = 1, U\right) = 2v_a\left(U/2\right) \tag{19}$$

for sufficiently large populations. Interestingly, Equation 19 is trivially valid in the SSWM regime where the speed is linear in U and $v(r, U) \approx 2NUs^2$ irrespective of r. Since Equation 19 accurately estimates the speed for very small and very large populations, it is likely that Equation 19 is a good approximation for any population size. Indeed, as we show in Fig. 4, $v_s(r = 1, U)$ is well approximated by twice $v_a(U/2)$ for any population size. The parameters we have used in these simulations are $U = 10^{-6}$ and $s = 0.01$. With the help of Equation 18, we may thus approximate the speed v_s of the obligately sexual population as

$$\ln N \approx \frac{v_s}{4s^2}\left(\ln^2\frac{v_s}{eUs} + 1\right) + \ln\frac{v_s}{2s^2U}. \tag{20}$$

[Figure 4 about here.]

It is clear that for $0 < r \ll 1$ there should be a regime where Equation 19 cannot approximate the speed accurately. To see this deviation, we simulated populations with various r (Fig. 4). It turns out that Equation 19 is still a good approximation for $r \geq 10^{-2}$. In particular, the speed for $r = 0.1$ is hardly discernible from that for $r = 1$ for all population sizes. The deviation starts to be significant for $r = 10^{-3}$. For comparison, we also plot

$v_s(r=0,U)/v_a(U/2)$ or equivalently $v_a(U)/v_a(U/2)$ in Fig. 4, which should approach to 1 in the infinite population limit (PARK *et al.* 2010). For $N \geq 10^6$, where $NU \ln(Ns)$ becomes larger than 1, $v_s(r,U)/v_a(U/2)$ starts to increase though very slowly and $v_s(r,U)$ becomes significantly larger than $v_s(r=0,U) = v_a(U)$. Note that for asexual populations clonal interference sets in around $NU \ln(Ns) \sim 1$ (WILKE 2004; PARK *et al.* 2010). That is, as soon as clonal interference becomes relevant, even a small amount of recombination leads to a significant speedup of adaptation, in agreement with the FM mechanism. The fact that the ratio v_s/v_a continues to rise monotonically with N for all cases with $r > 0$ in Fig. 4 is consistent with the two-fold advantage predicted by the infinite population analysis.

When measuring the speed of adaptation in our simulations, a useful consistency check was provided by the Guess relation

$$v = Us + \left\langle \frac{1}{N} \sum_{i=1}^{N} (\chi_i - 1) \ln \chi_i \right\rangle_{\text{stat}}, \tag{21}$$

where χ_i is the relative fitness of i-th individual in the infinite time limit,

$$\chi_i = \lim_{t\to\infty} \frac{w_i(t)}{\bar{w}(t)}, \tag{22}$$

and $\langle \cdot \rangle_{\text{stat}}$ signifies an average over the stationary measure of χ_i. Equation 21 was originally established for asexual populations undergoing discrete generation (WF) dynamics (GUESS 1974a; GUESS 1974b). In APPENDIX B, we prove that the relation holds for sexuals as well, and in Fig. 5, we numerically confirm its validity. The two terms on the right hand side of Equation 21 represent the increase in population fitness due to mutation and selection, respectively. Recombination affects the speed of adaptation only indirectly through its effect on the relative fitnesses χ_i. Note that the Guess relation should hold even if one uses discrete-time, overlapping generation models such as the Moran model.

[Figure 5 about here.]

Finally, we analyze the difference in the number of beneficial mutations acquired by the two loci. We quantify this difference as

$$\sigma \equiv \lim_{t\to\infty} \frac{\left\langle (\ell_1(t) - \ell_2(t))^2 \right\rangle}{t}, \tag{23}$$

where ℓ_1 and ℓ_2 are defined in Equation 16. We will refer to σ as the mutation number imbalance (MNI). To discern the MNI of asexuals from that of sexuals, we will add subscripts a and s, for asexual and sexual populations, respectively. In infinite populations each locus accumulates the same number of mutations, hence this study is meaningful only for finite populations.

For asexual populations, an approximation for σ can be obtained by comparing the *origination processes* at the two loci, which count the mutations that are present in some individuals of the population at time t and that are destined to eventually go to fixation (Gillespie 1993; Gillespie 1994; Park and Krug 2007). Denoting the number of such mutations at locus i by $k_i(t)$, we assume that 1) the difference between $k_i(t)$ and the lead $\ell_i(t) \geq k_i(t)$ remains bounded in the long time limit, 2) the total number of mutations $M_F = k_1 + k_2$ in the origination process increases at the same rate as the mean number of mutations, $M_F(t) \approx v_a t/s$ for large t, and 3) each new mutation appearing in the origination process chooses one of two loci with equal probability. Assumptions 1) and 2) reflect the existence of a steady state and have been verified in simulations (Park and Krug 2007), and assumption 3) is a consequence of the symmetry between the two loci. By assumption 3), the probability that there are m_1 mutations at locus 1 and $m_2 = M_F(t) - m_1$ mutations at locus 2 is given by

$$P(m_1, t) \approx \binom{M_F(t)}{m_1} \left(\frac{1}{2}\right)^{M_F(t)}. \tag{24}$$

Since the mean of m_1 is $\langle m_1 \rangle = M_F/2$ and its variance is $\langle (m_1 - \langle m_1 \rangle)^2 \rangle = M_F/4$, we can calculate σ_a, invoking the assumption 1), as

$$\sigma_a \approx \frac{\left\langle (M_F(t) - 2m_1)^2 \right\rangle}{t} = \frac{4}{t} \left\langle (m_1 - \langle m_1 \rangle)^2 \right\rangle \to \frac{v_a}{s}. \tag{25}$$

In Figure 6, we compare σ_a to v_a/s for $U = 10^{-6}$ and $s = 0.01$, which shows an excellent agreement.

[Figure 6 about here.]

Recombination changes the behavior of the MNI substantially. As can be seen in Figure 6, once the population size is in the regime of clonal interference the MNI decreases abruptly, then remains almost constant for a wide range of population sizes. Even a small amount of recombination efficiently equalizes any major fitness difference between the two loci by creating competitively superior recombinants in which both loci have high fitness.

DISCUSSION

The Fisher-Muller mechanism for the evolutionary advantage of sex is based on the slowing down of asexual adaptation due to clonal interference, which is alleviated by the recombination of high fitness genotypes. While much recent theoretical work has been devoted to quantifying the speed of adaptation in asexuals, the speedup that can be achieved through recombination has been explicitly addressed only in a few studies (see below). In the present article we take a step in this direction by providing a detailed analysis of a simple, yet biologically meaningful model in which recombination occurs between two loci, each of which can harbor an unlimited number of linked beneficial mutations. Our analysis shows that the advantage of sex becomes significant in the parameter regime where clonal interference plays an important role in asexual populations. In our two-locus model, the adaptation speed of sexual populations is about twice as large as that of the corresponding asexual populations for a wide range of recombination rate, an effect that we refer to as a two-fold advantage of sex. In the remainder of this section we discuss the extension of our model to L loci and compare our findings to related previous work.

More than two loci: It is natural to surmise that the factor of two arises in our model because we are considering two loci, and that the speed increase should in general by proportional to the number of loci. Indeed, this turns out to be true if we use a ‘communal’ recombination scheme where the gene of each locus is collected from the whole population rather than from the two parents, once the genome of the offspring is constructed by recombination. In a three-locus model with this recombination scheme, the frequency distribution of next generation is sampled from

$$f_t^r(n_1, n_2, n_3) = (1-r) f_t^\mu(n_1, n_2, n_3) + r f_t^{(1)}(n_1) f_t^{(2)}(n_2) f_t^{(3)}(n_3), \tag{26}$$

where $f_t^{(i)}(n_i)$ is the marginal frequency distribution for having n_i mutations at locus i after the (deterministic) selection and mutation steps (compare to MODELS). It is a straightforward extension of the calculation in APPENDIX A to show that the infinite population dynamics for $r = 1$ is again divided into three independent evolutions of each locus with marginal mutation probabilities just as in the two-locus case. In general, if we consider a model system with L loci within the communal recombination scheme mentioned above, the evolution is a superposition of L independent evolutions at each locus, and there is an L-fold advantage of sex in the infinite population.

To see if this L-fold advantage persists for finite populations, we performed simulations of the three-locus model. As in the two-locus case, we expect that

$$v_s(r, U) = 3 v_a(U/3) \tag{27}$$

for sufficiently large r. Indeed, we observe that the simulations are consistent with Equation 27 for a wide range of parameter values. In Fig. 7, we depict $v_s(r, U)/v_a(U/3)$ as a function of N for $U = 1.5 \times 10^{-6}$ and $s = 0.01$ with varying r. As in the two-locus model, the advantage of sex becomes significant when clonal interference is important in the corresponding asexual populations.

[Figure 7 about here.]

We also studied the mutation number inbalance in the three-locus model. We slightly modify the definition of the MNI as the difference between the maximum and minimum numbers of accumulated mutations at all loci, which reduces to the definition of Equation 23 for the two-locus model. In Fig. 8, we depict the MNI for the three-locus model with $U = 1.5 \times 10^{-7}$ and $s = 0.01$ for various r. Like the MNI of the two-locus model, the MNI for the asexuals increases with N while slowly decreasing for sexuals. Again, the qualitative difference between sexuals and asexuals becomes significant in the regime where clonal intereference is important.

[Figure 8 about here.]

Relation to previous studies: The quantitative analysis of the speed of evolution of sexual populations compared to that of asexual populations, when both evolve on the same non-epistatic fitness landscape with the same beneficial mutation rate per genome, has a long history (Crow and Kimura 1965; Maynard Smith 1968; Crow and Kimura 1969; Maynard Smith 19 Felsenstein 1974; Maynard Smith 1978; Kim and Orr 2005). When the number of accessible beneficial mutations is finite, the relevant quantity is the time for all beneficial mutations to be fixed. In this context, Maynard Smith (1971) analyzed the fixation time for sexual and asexual populations evolving on a fitness landscape with l loci under selection. Each locus has two alleles, one of which confers a beneficial fitness effect in a non-epistatic fashion. For sexual populations, the linkage among loci was assumed weak. Using a rough approximation, Maynard Smith (1971) argued that the time for completing evolutionary changes in asexual populations is l times longer than that in sexual populations for sufficiently large population size, which implies an l-fold advantage of sex similar to what we found in our study. While the analysis by Maynard Smith (1971) is fairly crude and (as conceded by the author) actually not consistent with the simulation results presented in the same paper, the conclusion that the advantage of sex becomes stronger with an increasing number of loci under selection is in qualitative agreement with our results, as well as with the related work of Kim and Orr (2005).

Recent studies of the speed of sexual populations in the context of the FM mechanism have mostly focused on the case with an infinite supply of beneficial mutations (Neher *et al.* 2010;

Rouzine and Coffin 2010; Weissman and Barton 2012), exploiting the mathematical progress in treating the spreading of beneficial mutations as a Gaussian traveling wave (Tsimring *et al.* 199 Rouzine *et al.* 2003; Desai and Fisher 2007; Rouzine *et al.* 2008; Park *et al.* 2010; Good *et al.* 2012; Fisher 2013). Rouzine and Coffin (2010) studied how recombination speeds up adaptation when there is standing variation of beneficial mutations. Neher *et al.* (2010) studied the speed of adaptation of large facultatively sexual populations, starting from a monomorphic state. Similar to our results, Neher *et al.* (2010) found a regime of intermediate recombination rates where the speed increases logarithmically with population size, however with a prefactor that varies quadratically with r. Although Rouzine and Coffin (2010) and Neher *et al.* (2010) investigated the adaptation dynamics of sexual populations with an (effectively) infinite supply of beneficial mutations, their results cannot be directly compared to ours. This is because the model genomes of Rouzine and Coffin (2010) and Neher *et al.* (2010) assume no or weak linkage between beneficial mutations, whereas in our model mutations in the same locus are tightly linked. A related study with an explicit genetic map was recently presented by Weissman and Barton (2012).

The two-locus genome considered in this paper can be viewed as a simple example of a modular genomic architecture, where recombination occurs between modules but not within a modul. Watson *et al.* (2011) have pointed out that such a modular structure induces a strong benefit for sexual reproduction when there is sign epistasis within the modules and different modules contribute independently to fitness. It would therefore be of interest to extend our approach to include a tunable degree of epistasis interactions within the loci. Following Watson *et al.* (2011), such interactions should affect not only the speed of adaptation but also the set of genotypes that can be reached at all by the population.

ACKNOWLEDGMENTS

Support by Deutsche Forschungsgemeinschaft within SFB 680 *Molecular Basis of Evolutionary Innovations* is gratefully acknowledged. In addition, S.-C.P. acknowledges the support by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (Grant No. 2011-0014680) and the Catholic University of Korea, Research Fund, 2013.

LITERATURE CITED

Barrick, J. E., D. S. Yu, S. H. Yoon, H. Jeong, T. K. Oh, D. Schneider, R. E. Lenski, and J. F. Kim, 2009 Genome evolution and adaptation in a long-term experiment with *Escherichia coli*. Nature **461:** 1243–1247.

Brunet, E., I. M. Rouzine, and C. O. Wilke, 2008 The stochastic edge in adaptive evolution. Genetics **179:** 603–620.

Charlesworth, D., M. Morgan, and B. Charlesworth, 1993 Mutation accumulation in finite outbreeding and in- breeding populations. Genet. Res. Camb. **61:** 39–56.

Colegrave, N., 2002 Sex releases the speed limit on evolution. Nature **420:** 664–666.

Cooper, T. F., 2007 Recombination speeds adaptation by reducing competition between beneficial mutations in populations of *Escherichia coli*. PLoS Biol. **5:** e225.

Crow, J. and M. Kimura, 1969 Evolution in sexual and asexual populations: A reply. Am. Nat. **103:** 89–91.

Crow, J. F. and M. Kimura, 1965 Evolution in sexual and asexual populations. Am. Nat. **99:** 439–450.

de Visser, J. A. G. M. and S. F. Elena, 2007 The evolution of sex: empirical insights into the roles of epistasis and drift. Nature Reviews Genetics **8:** 139–149.

de Visser, J. A. G. M., S.-C. Park, and J. Krug, 2009 Exploring the effect of sex on empirical fitness landscapes. Am. Nat. **174:** S15–S30.

Desai, M. M. and D. S. Fisher, 2007 Beneficial mutation-selection balance and the effect of linkage on positive selection. Genetics **176:** 1759–1798.

Eshel, I. and M. Feldman, 1970 On the evolutionary effect of recombination. Theor. Popul. Biol. **1:** 88–100.

Felsenstein, J., 1974 The evolutionary advantage of recombination. Genetics **78:** 737–756.

Fisher, D. S., 2013 Asexual evolution waves: fluctuations and universality. J. Stat. Mech.:Theory Exp.**:** P01011.

Fisher, R. A., 1930 *The Genetical Theory of Natural Selection*. Clarendon Press, Oxford.

Fogle, C. A., J. L. Nagle, and M. M. Desai, 2008 Clonal interference, multiple mutations and adaptation in large asexual populations. Genetics **180:** 2163–2170.

Franke, J., A. Klözer, J. A. G. M. de Visser, and J. Krug, 2011 Evolutionary accessibility of mutational pathways. PLoS Comp. Biol. **7:** e1002134.

Gerrish, P. J., 2001 The rhythm of microbial adaptation. Nature **413:** 299–302.

Gerrish, P. J. and R. E. Lenski, 1998 The fate of competing beneficial mutations in an asexual population. Genetica **102-103:** 127–144.

Gillespie, J. H., 1993 Substitution processes in molecular evolution: I. uniform and clustered substitutions in a haploid model. Genetics **134:** 971–981.

Gillespie, J. H., 1994 *The Causes of Molecular Evolution*. Oxford University Press, Oxford.

Good, B. H., I. M. Rouzine, D. J. Balick, O. Hallatschek, and M. M. Desai, 2012 Distribution of fixed beneficial mutations and the rate of adaptation in asexual populations. Proc. Nat. Acad. Sci. USA **109:** 4950–4955.

Guess, H. A., 1974a Evolution in finite population with infinitely many types. Theor. Popul. Biol. **5:** 417–430.

Guess, H. A., 1974b Limit theorems for some stochastic evolution models. Ann. Prob. **2:** 14–31.

Jain, K., 2008 Loss of least-Loaded class in asexual populations due to drift and epistasis. Genetics **179:** 2125–2134.

Jain, K., S.-C. Park, and J. Krug, 2011 Evolutionary advantage of small populations on complex fitness landscapes. Evolution **65:** 1945 – 1955.

Kim, Y. and H. A. Orr, 2005 Adaptation in sexual *vs.* asexuals: clonal interference and the Fisher-Muller model. Genetics **171:** 1377–1386.

Kimura, M., 1969 The number of heterozygous nucleotide sites maintained in a finite population due to steady flux of mutations. Genetics **61:** 893–903.

Kingman, J. F. C., 1978 A simple model for the balance between selection and mutation. J. Appl. Prob. **15:** 1–12.

Kondrashov, A. S., 1988 Deleterious mutations and the evolution of sexual reproduction. Nature **336:** 435–440.

Kondrashov, A. S., 1993 Classification of hypotheses on the advantage of amphimixis. J. Hered. **84:** 372–387.

Kondrashov, A. S., 1994 Muller's ratchet under epistatic selection. Genetics **136:** 1469–1473.

Kouyos, R. D., G. E. Leventhal, T. Hinkley, M. Haddad, J. M. Whitcomb, C. J. Petropoulos, and S. Bonhoeffer, 2012 Exploring the Complexity of the HIV-1 Fitness Landscape. PLoS Genet. **8:** e1002551.

Lenski, R. E., M. R. Rose, S. C. Simpson, and S. C. Tadler, 1991 Long-term experimental evolution in *Escherichia coli.* I. adaptation and divergence during 2,000 generations. Am. Nat. **138:** 1315–1341.

Lenski, R. E. and M. Travisano, 1994 Dynamics of adaptation and diversification: A 10,000-generation experiment with bacterial populations. Proc. Nat. Acad. Sci. USA **91:** 6808–6814.

Maynard Smith, J., 1968 Evolution in sexual and asexual populations. Am. Nat. **102:** 469–473.

Maynard Smith, J., 1971 What use is sex? J. Theor. Biol. **30:** 319–335.

Maynard Smith, J., 1978 *The evolution of sex*. Cambridge University Press, Cambridge.

Miralles, R., P. J. Gerrish, A. Moya, and S. F. Elena, 1999 Clonal interference and the evolution of RNA viruses. Science **285:** 1745–1747.

Muller, H. J., 1932 Some genetic aspects of sex. Am. Nat. **66:** 118–138.

Muller, H. J., 1964 The relation of recombination to mutational advance. Mutat. Res. **1:** 2–9.

Mustonen, V. and M. Lässig, 2010 Fitness flux and ubiquity of adaptive evolution. Proc. Nat. Acad. Sci. USA **107:** 4248–4253.

Neher, R. A., B. I. Shraiman, and D. S. Fisher, 2010 Rate of adaptation in large sexual populations. Genetics **184:** 467–481.

Otto, S. P., 2009 The evolutionary enigma of sex. Am. Nat. **174:** S1–S14.

Park, S.-C. and J. Krug, 2007 Clonal interference in large populations. Proc. Nat. Acad. Sci. USA **104:** 18135–18140.

Park, S.-C. and J. Krug, 2008 Evolution in random fitness landscapes: the infinite sites model. J. Stat. Mech. **2008:** P04014.

Park, S.-C. and J. Krug, 2011 Bistability in two-locus models with selection, mutation, and recombination. J. Math. Biol. **62:** 763–788.

Park, S.-C., D. Simon, and J. Krug, 2010 The speed of evolution in large asexual populations. J. Stat. Phys. **138:** 381–410.

Rouzine, I. M., E. Brunet, and C. O. Wilke, 2008 The traveling-wave approach to asexual evolution: Muller's ratchet and speed of adaptation. Theor. Popul. Biol. **73:** 24–46.

Rouzine, I. M. and J. M. Coffin, 2010 Multi-site adaptation in the presense of infrequent recombination. Theor. Popul. Biol. **77:** 189–204.

Rouzine, I. M., J. Wakeley, and J. M. Coffin, 2003 The solitary wave of asexual evolution. Proc. Nat. Acad. Sci. USA **100:** 587–592.

Schiffels, S., G. J. Szöllösi, V. Mustonen, and M. Lässig, 2011 Emergent Neutrality in Adaptive Asexual Evolution. Genetics **189:** 1361–1375.

Sniegowski, P. D. and P. J. Gerrish, 2010 Beneficial mutations and the dynamics of adaptation in asexual populations. Proc. R. Soc. Lond. Ser. B **365:** 1255–1263.

Szendro, I. G., J. Franke, J. A. G. de Visser, and J. Krug, 2013 Predictability of evolution depends nonmonotonically on population size. Proc. Nat. Acad. Sci. USA **110:** 571–576.

Szendro, I. G., M. F. Schenk, J. Franke, J. Krug, and J. A. G. de Visser, 2013 Quantitative analyses of empirical fitness landscapes. J. Stat. Mech.:Theory Exp.**:** P01005.

Tsimring, L. S., H. Levine, and D. A. Kessler, 1996 RNA virus evolution via a fitness-space model. Phys. Rev. Lett. **76:** 4440–4443.

Watson, R. A., D. M. Weinreich, and J. Wakeley, 2011 Genome structure and the benefit of sex. Evolution **65:** 523–536.

Weinreich, D. M., N. F. Delaney, M. A. DePristo, and D. L. Hartl, 2006 Darwinian evolution can follow only very few mutational paths to fitter proteins. Science **312:** 111–114.

Weinreich, D. M., R. A. Watson, and L. Chao, 2005 Sign epistasis and genetic constraint on evolutionary trajectories. Evolution **59:** 1165–1174.

Weissman, D. B. and N. H. Barton, 2012 Limits to the rate of adaptive substitution in sexual populations. PLoS Genet. **8:** e1002740.

Wilke, C. O., 2004 The Speed of Adaptation in Large Asexual Populations. Genetics **167:** 2045–2053.

Wright, S., 1931 Evolution in Mendelian populations. Genetics **16:** 97–159.

APPENDIX A: INFINITE POPULATION DYNAMICS FOR ASEXUALS ($r = 0$) AND OBLIGATE SEXUALS ($r = 1$)

Let $F_t(z_1, z_2)$ denote the generating function for the frequency distribution at generation t, which is defined as

$$F_t(z_1, z_2) \equiv \sum_{n_1,n_2} z_1^{n_1} z_2^{n_2} f_t(n_1, n_2). \tag{A1}$$

Since the fitness landscape is multiplicative, the mean fitness at generation t can be found from F_t through

$$\bar{w}_t = F_t(e^{s_1}, e^{s_2}). \tag{A2}$$

Likewise, we introduce the generating function for f_t^s in Equation 1, which is obtained from F_t according to

$$F_t^s(z_1, z_2) \equiv \sum_{n_1,n_2} z_1^{n_1} z_2^{n_2} f_t^s(n_1, n_2) = \frac{F_t(e^{s_1} z_1, e^{s_2} z_2)}{F_t(e^{s_1}, e^{s_2})}. \tag{A3}$$

Since f_t^μ in Equation 3 is the convolution of g_0 and f_t^s, the generating function for f_t^μ is the product of F_t^s and $G_0(z_1, z_2)$ (for the definition of the latter quantity see Equation 10).

Using that f_{t+1} is the same as f_t^r for infinite populations, we obtain an iterative evolution equation for F_t that reads

$$\begin{aligned} F_{t+1}(z_1, z_2) &= (1-r)F_t^\mu(z_1, z_2) + rF_t^\mu(1, z_2)F_t^\mu(z_1, 1) & \text{(A4)} \\ &= (1-r)G_0(z_1, z_2)\frac{F_t(z_1 e^{s_1}, z_2 e^{s_2})}{F_t(e^{s_1}, e^{s_2})} + r\widetilde{G}_0(z_1)\frac{F_t(z_1 e^{s_1}, e^{s_2})}{F_t(e^{s_1}, e^{s_2})}\widetilde{G}_0(z_2)\frac{F_t(e^{s_1}, z_2 e^{s_2})}{F_t(e^{s_1}, e^{s_2})}, \end{aligned}$$

where $F_t^\mu(z_1, z_2) = G_0(z_1, z_2)F_t^s(z_1, z_2)$ is the generating function of f_t^μ and

$$\widetilde{G}_0(z) = \sum_{k_1,k_2} z^{k_1} g_0(k_1, k_2) = \sum_{k_1,k_2} z^{k_2} g_0(k_1, k_2). \tag{A5}$$

For $r = 0$, Equation A4 can be solved by iterating backwards until $t = 0$, that is (Park and Krug 2007)

$$F_t(z_1, z_2) = G_0(z_1, z_2)\frac{F_{t-1}(z_1 e^{s_1}, z_2 e^{s_2})}{F_{t-1}(e^{s_1}, e^{s_2})} = G_0(z_1, z_2)\frac{G_0(z_1 e^{s_1}, z_2 e^{s_2})}{G_0(e^{s_1}, e^{s_2})}\frac{F_{t-2}(z_1 e^{2s_1}, z_2 e^{2s_2})}{F_{t-2}(e^{2s_1}, e^{2s_2})},$$

$$= \frac{F_0(z_1 e^{ts_1}, z_2 e^{ts_2})}{F_0(e^{ts_1}, e^{ts_2})}\prod_{\tau=0}^{t-1}\frac{G_0(z_1 e^{s_1\tau}, z_2 e^{s_2\tau})}{G_0(e^{s_1\tau}, e^{s_2\tau})} \quad \text{(A6)}$$

with $F_0(z_1, z_2) = 1$ for the homogeneous initial condition. Thus the mean fitness at generation t is

$$\bar{w}_t = G_0(e^{s_1 t}, e^{s_2 t}). \quad \text{(A7)}$$

When $r = 1$, Equation A4 suggests that the generating function is the product of two functions such that

$$F_t(z_1, z_2) = \widetilde{F}_t^1(z_1)\widetilde{F}_t^2(z_2). \quad \text{(A8)}$$

With the above ansatz, we can find an evolution equation for $\widetilde{F}_t^i(z)$ $(i = 1 \text{ or } 2)$ from Equation A4,

$$\widetilde{F}_{t+1}^i(z) = \widetilde{G}_0(z)\frac{\widetilde{F}_t^i(ze^{s_i})}{\widetilde{F}_t^i(e^{s_i})}, \quad \text{(A9)}$$

which is exactly the evolution equation for an asexual population with mutation generating function $\tilde{G}_0(z)$. The solution of Equation A9 is

$$\widetilde{F}_t^i(z) = \prod_{\tau=0}^{t-1}\frac{\widetilde{G}_0(ze^{s_i\tau})}{\widetilde{G}_0(e^{s_i\tau})}, \quad \text{(A10)}$$

where we have used the initial condition $F_0(z_1, z_2) = \widetilde{F}_0(z) = 1$. One can easily check that Equation A8 with $\tilde{F}_t(z)$ given by Equation A9 actually solves Equation A4 for $r = 1$ by substitution. Hence the mean fitness at generation t for $r = 1$ is

$$w_t = F_t(e^{s_1}, e^{s_2}) = \widetilde{G}_0(e^{s_1 t})\widetilde{G}_0(e^{s_2 t}). \quad \text{(A11)}$$

APPENDIX B: GUESS RELATION IN THE PRESENCE OF RECOMBINATION

In this APPENDIX, we will show that for evolution on multiplicative, non-epistatic fitness landscapes the Guess relation (Equation 21) is valid even in the presence of recombination.

Let $w_i(t)$ be the fitness of the i-th individual at generation t, $\bar{w}(t)$ the mean fitness of the population, $\bar{w}(t) = \sum_i w_i(t)/N$, and $X_i(t)$ the relative fitness of i-th individual, $X_i(t) = w_i(t)/\bar{w}(t)$. We will assume that $X_i(t)$ approaches a well-defined steady state as t goes to infinity. We take each individual to be characterized by a genome with L loci, each of which has inifitely many sites. The contribution of a locus to fitness is denoted by z_n $(n = 1, \ldots, L)$ and the fitness of an individual with such a genome is $w = \prod_{n=1}^{L} z_n$. In the following, z_n will be called CFn, meaning the Contribution to Fitness of the nth-locus. If a mutation hits the n-th locus, CFn changes from z_n to $z'_n = z_n v_n$, where v_n is drawn from a given probability distribution that may vary from locus to locus but does not depend on z_n or the generation. If v_n is larger (smaller) than 1, the mutation is beneficial (deleterious). In this APPENDIX, the explicit form of the probability distribution for v_n does not need to be specified.

We will use the vector notation $\vec{z} = (z_1, ..., z_L)$ for fitness vectors with L elements. Assume that there are N individuals and the CFn of individual i is $z_{i,n}$. The corresponding fitness vector is denoted by $\vec{z}_i$. At first, we will calculate the expected mean log-fitness at generation $t + 1$ assuming that $X_i(t)$ and $\vec{z}_i(t)$ are given.

By selection, the probability density that the fitness vector of an offspring is $\vec{z}$ is

$$f_s(\vec{z}) = \sum_{i=1}^{N} \frac{X_i(t)}{N} \delta(\vec{z} - \vec{z}_i), \tag{B1}$$

where $\delta(\vec{x})$ is the L-dimensional Dirac delta function. Let $g(\vec{v})$ be the probability density that a mutation event changes the CFn of an offspring by v_n $(z_n \to z'_n = z_n v_n)$ for all n's. Then due to mutation, the expected frequency becomes

$$f_m(\vec{z}) = \int d\vec{v} d\vec{z}' \delta(\vec{z} - \vec{z}' \otimes \vec{v}) g(\vec{v}) f_s(\vec{z}'), \tag{B2}$$

where $\vec{z}' \otimes \vec{v}$ denotes the vector with elements $z'_n v_n$.

Next we consider recombination. Let $R(\vec{z}|\vec{z}_1, \vec{z}_2)$ be the probability density that offspring resulting from the recombination of two parents with fitness vectors $\vec{z}_1$ and $\vec{z}_2$ has fitness $\vec{z}$.

In general, we can write R in the form

$$R(\vec{z}|\vec{z}_1, \vec{z}_2) = \sum_S p(S) \prod_{n=1}^{L} \delta\left(z_n - z_{S(n),n}\right), \tag{B3}$$

where S runs over all possible outcomes of recombination and $p(S)$ is the probability of this event. Here $S(n) = 1$ (2) if locus n is inherited from parent 1 (2), hence the total number of possible outcomes is 2^L. Then the final probability density becomes

$$f(\vec{z}) = \int d\vec{z}_1 d\vec{z}_2 R(\vec{z}|\vec{z}_1, \vec{z}_2) f_m(\vec{z}_1) f_m(\vec{z}_2). \tag{B4}$$

Now we will calculate the expected log-fitness of a randomly chosen individual at generation $t+1$ for given $w_i(t)$. Since the probability density that a randomly chosen individual at generation $t+1$ has fitness vector $\vec{z}$ is $f(\vec{z})$ given in Equation B4 and the corresponding fitness is $w = \prod_n z_n$, the quantity we want to calculate is

$$I \equiv \int dw \ln w \ \text{prob}(w) = \int d\vec{z} \ln\left(\prod_{n=1}^{L} z_n\right) f(\vec{z}), \tag{B5}$$

where prob(w) is the probability density that an individual has fitness w at generation $t+1$ for given $w_i(t)$. Guess (1974b) showed that as long as there is a well-defined steady state the speed v can be calculated as

$$v = \left\langle \ln \frac{\bar{w}(t+1)}{\bar{w}(t)} \right\rangle = \left\langle \ln \frac{\mathcal{W}_1}{\mathcal{W}_2} \right\rangle = \langle I \rangle - \langle \ln \mathcal{W}_2 \rangle, \tag{B6}$$

where $\mathcal{W}_1$ ($\mathcal{W}_2$) is the fitness of a randomly chosen individual at generation $t+1$ (t) and $\langle \ldots \rangle$ signifies the average over the steady state distribution. Loosely speaking, the above relation can be understood as follows: Since the dynamics is symmetric under permutations of the population index, the steady state must have this permutation symmetry as well. Hence the expected log-mean fitness at steady state should be the same as the expected log-fitness of a randomly chosen individual. Moreover, at stationarity the speed can be calculated from the difference of log-fitness between two consecutive generations. Thus Equation B6 follows.

From Equations B1, B2, B3, and B4, we get

$$\begin{aligned}
I &= \int d\vec{z} \ln\left(\prod_{n=1}^{L} z_n\right) \sum_{i,j} \frac{X_i(t)X_j(t)}{N^2} \int d\vec{v}_i d\vec{v}_j R(\vec{z}|\vec{z}_i \otimes \vec{v}_i, \vec{z}_j \otimes \vec{v}_j) g(\vec{v}_i) g(\vec{v}_j) \\
&= \sum_{i,j} \frac{X_i(t)X_j(t)}{N^2} \int d\vec{v}_i d\vec{v}_j g(\vec{v}_i) g(\vec{v}_j) \int d\vec{z} \ln\left(\prod_{n=1}^{L} z_n\right) R(\vec{z}|\vec{z}_i \otimes \vec{v}_i, \vec{z}_j \otimes \vec{v}_j) \\
&= \sum_{i,j} \sum_{S} p(S) \frac{X_i(t)X_j(t)}{N^2} \int d\vec{v}_i d\vec{v}_j g(\vec{v}_i) g(\vec{v}_j) \sum_{n=1}^{L} \left[\ln(z_{S(n),n}) + \ln(v_{S(n),n})\right] \\
&= \sum_{S} p(S) \sum_{n=1}^{L} \left(\sum_{i,j} \frac{w_i w_j}{N^2} \ln(z_{S(n),n}) + \int d\vec{v}_i d\vec{v}_j g(\vec{v}_i) g(\vec{v}_j) \ln(v_{S(n),n}) \right), \qquad \text{(B7)}
\end{aligned}$$

where $S(n)$ in the subscript of z and v can be either i or j. Since

$$\sum_{ij} \frac{X_i(t)X_j(t)}{N^2} \ln(z_{i,n}) = \sum_{ij} \frac{X_i(t)X_j(t)}{N^2} \ln(z_{j,n}) = \sum_{i} \frac{X_i(t)}{N} \ln(z_{i,n}), \qquad \text{(B8)}$$

$$\int d\vec{v}_i d\vec{v}_j g(\vec{v}_i) g(\vec{v}_j) \ln v_{i,n} = \int d\vec{v}_i d\vec{v}_j g(\vec{v}_i) g(\vec{v}_j) \ln v_{j,n} = \int d\vec{v} g(\vec{v}) \ln v_n, \qquad \text{(B9)}$$

the summation and integral in the parentheses of the last line of Equation B7 do not depend on S. Due to the normalization $\sum_S p(S) = 1$, we get

$$I = \sum_i \frac{X_i(t)}{N} \ln(X_i(t)\bar{w}(t)) + \langle \ln V \rangle = \sum_i \frac{X_i(t)}{N} \ln(X_i(t)) + \langle \ln V \rangle + \ln \bar{w}(t), \qquad \text{(B10)}$$

where V is the total effect of fitness increase by a single mutation event, $V = \prod_n v_n$, and $\prod_n z_{i,n} = X_i(t)\bar{w}(t)$ was used. Hence the speed becomes

$$v = \langle I \rangle - \left\langle \sum_i \frac{1}{N} \ln w_i \right\rangle = \langle \ln V \rangle + \left\langle \frac{1}{N} \sum_i (\chi_i - 1) \ln \chi_i \right\rangle, \qquad \text{(B11)}$$

where χ_i is the relative fitness of the individual i at steady state. Note that the formula does not depend on the explicit form of the recombination operator. If we use Equation 15 for the mutation scheme, $\langle \ln V \rangle = Us$.

The application of the above procedure to the Moran model is straightforward. Hence, the Guess relation is valid for discrete time models regardless of recombination, once the fitness landscape is multiplicative.

List of Figures

1 Mean fitness $\bar{w}_t$ of the infinite population model as a function of time for $r = 0$, $r = 10^{-9}$, and $r = 1$ (from bottom to top) with $U = 0.1$ and $s = 0.02$. As argued in the text, the speed does not depend on r once r is nonzero. . . 34

2 Frequency distribution of the total number of mutations for the infinite population model at generations $895, 900, 905, \ldots, 930$ (left to right) with parameters $r = 10^{-9}$, $U = 0.1$, and $s = 0.02$. 35

3 Speed of adaptation of finite asexual populations as a function of N on a double logarithmic scale for $U = 10^{-6}$ and $s = 0.01$. The numerical solutions of Equation 17 and Equation 18 are drawn for comparison with the simulation data. As anticipated, the *ad-hoc* modification (Equation 18) provides a more accurate estimate. 36

4 Ratio of the sexual adaptation speed, $v_s(r, U)$, to the asexual speed at half mutation rate, $v_a(U/2)$, as a function of population size N, where v_s is obtained from simulations and v_a is computed using Equation 18. Recombination rates are $r = 0$ (empty reverse triangle), 10^{-5} (empty square), 10^{-4} (filled triangle), 10^{-3} (empty triangle),10^{-2} (filled circle),10^{-1} (empty circle), and 1 (filled square) from bottom to top, and $U = 10^{-6}$ and $s = 0.01$ are used throughout. The scaling relation in Equation 19 predicts that $v_s(r, U)/v_a(U/2) = 2$. Note that two datasets for $r = 0.1$ (empty circle) and $r = 1$ (filled square) are indiscernible. 37

5 Numerical verification of the Guess relation. The figure compares the logarithmic mean fitness divided by generation time, $\langle \ln \bar{w}(t) \rangle / t$, to the right hand side of Equation 21. The data are obtained from simulations with $r = 10^{-3}$, $U = 10^{-6}$ and $s = 0.01$. For $t \to \infty$, both curves intersect the ordinate at the same point, which equals the asymptotic speed of adaptation. 38

6 The mutation number imbalance (MNI) σ vs. population size N for $r = 0$ (empty reverse triangle), 10^{-5} (empty square), 10^{-4} (filled triangle), 10^{-3} (empty triangle),10^{-2} (filled circle),10^{-1} (empty circle), and 1 (filled square) from top to bottom. Other parameter values are $U = 10^{-6}$ and $s = 0.01$. The symbols for $r \geq 10^{-2}$ essentially overlap. For comparison, the analytic prediction $\sigma_a = v_a/s$ for asexuals is drawn as a line. 39

7 Ratio of the sexual adaptation speed in the three-locus model, $v_s(r, U)$, to the asexual speed v_a at mutation rate $U/3$ as a function of population size N, where v_s is obtained from simulations and v_a is computed using Equation 18. Recombination rates are $r = 0$ (empty reverse triangle), 10^{-5} (empty square), 10^{-4} (filled triangle), 10^{-3} (empty triangle),10^{-2} (filled circle),10^{-1} (empty circle), and 1 (filled square) from bottom to top, and $U = 10^{-6}$ and $s = 0.01$ are used throughout. The scaling relation in Equation 27 predicts that $v_s(r, U)/v_a(U/3) = 3$. Note that two datasets for $r = 0.1$ (empty circle) and $r = 1$ (filled square) are indiscernible. 40

8 Mutation number imbalance (MNI) for the three-locus model with recombination rates $r = 0$ (empty reverse triangle), 10^{-5} (empty square), 10^{-4} (filled triangle), 10^{-3} (empty triangle),10^{-2} (filled circle),10^{-1} (empty circle), and 1 (filled square) from top to bottom. Other parameter values are $U = 1.5 \times 10^{-6}$ and $s = 0.01$. The symbols for $r \geq 10^{-2}$ essentially overlap. 41



Figure 1: Mean fitness $\bar{w}_t$ of the infinite population model as a function of time for $r = 0$, $r = 10^{-9}$, and $r = 1$ (from bottom to top) with $U = 0.1$ and $s = 0.02$. As argued in the text, the speed does not depend on r once r is nonzero.



Figure 2: Frequency distribution of the total number of mutations for the infinite population model at generations $895, 900, 905, \ldots, 930$ (left to right) with parameters $r = 10^{-9}$, $U = 0.1$, and $s = 0.02$.



Figure 3: Speed of adaptation of finite asexual populations as a function of N on a double logarithmic scale for $U = 10^{-6}$ and $s = 0.01$. The numerical solutions of Equation 17 and Equation 18 are drawn for comparison with the simulation data. As anticipated, the *ad-hoc* modification (Equation 18) provides a more accurate estimate.



Figure 4: Ratio of the sexual adaptation speed, $v_s(r,U)$, to the asexual speed at half mutation rate, $v_a(U/2)$, as a function of population size N, where v_s is obtained from simulations and v_a is computed using Equation 18. Recombination rates are $r = 0$ (empty reverse triangle), 10^{-5} (empty square), 10^{-4} (filled triangle), 10^{-3} (empty triangle),10^{-2} (filled circle),10^{-1} (empty circle), and 1 (filled square) from bottom to top, and $U = 10^{-6}$ and $s = 0.01$ are used throughout. The scaling relation in Equation 19 predicts that $v_s(r,U)/v_a(U/2) = 2$. Note that two datasets for $r = 0.1$ (empty circle) and $r = 1$ (filled square) are indiscernible.



Figure 5: Numerical verification of the Guess relation. The figure compares the logarithmic mean fitness divided by generation time, $\langle \ln \bar{w}(t) \rangle / t$, to the right hand side of Equation 21. The data are obtained from simulations with $r = 10^{-3}$, $U = 10^{-6}$ and $s = 0.01$. For $t \to \infty$, both curves intersect the ordinate at the same point, which equals the asymptotic speed of adaptation.



Figure 6: The mutation number imbalance (MNI) σ vs. population size N for $r = 0$ (empty reverse triangle), 10^{-5} (empty square), 10^{-4} (filled triangle), 10^{-3} (empty triangle),10^{-2} (filled circle),10^{-1} (empty circle), and 1 (filled square) from top to bottom. Other parameter values are $U = 10^{-6}$ and $s = 0.01$. The symbols for $r \geq 10^{-2}$ essentially overlap. For comparison, the analytic prediction $\sigma_a = v_a/s$ for asexuals is drawn as a line.



Figure 7: Ratio of the sexual adaptation speed in the three-locus model, $v_s(r, U)$, to the asexual speed v_a at mutation rate $U/3$ as a function of population size N, where v_s is obtained from simulations and v_a is computed using Equation 18. Recombination rates are $r = 0$ (empty reverse triangle), 10^{-5} (empty square), 10^{-4} (filled triangle), 10^{-3} (empty triangle),10^{-2} (filled circle),10^{-1} (empty circle), and 1 (filled square) from bottom to top, and $U = 10^{-6}$ and $s = 0.01$ are used throughout. The scaling relation in Equation 27 predicts that $v_s(r, U)/v_a(U/3) = 3$. Note that two datasets for $r = 0.1$ (empty circle) and $r = 1$ (filled square) are indiscernible.



Figure 8: Mutation number imbalance (MNI) for the three-locus model with recombination rates $r = 0$ (empty reverse triangle), 10^{-5} (empty square), 10^{-4} (filled triangle), 10^{-3} (empty triangle),10^{-2} (filled circle),10^{-1} (empty circle), and 1 (filled square) from top to bottom. Other parameter values are $U = 1.5 \times 10^{-6}$ and $s = 0.01$. The symbols for $r \geq 10^{-2}$ essentially overlap.